

FREE ENERGY IN THE HOPFIELD MODEL WITH MANY PATTERNS

JAMES Y. ZHAO

Department of Mathematics, Stanford University

Email: jyzhao@math.stanford.edu

12 September 2011

Abstract

We prove that in the Hopfield model where the ratio of patterns to sites grows large, the free energy behaves like the free energy in the Sherrington-Kirkpatrick model.

1 Introduction

The Hopfield model is the system of *configurations* $x \in \{\pm 1\}^N$ governed by the *Gibbs measure*

$$G(x) = \frac{1}{Z(\beta, B)} e^{-\beta H(x) + B \sum_i x_i}, \quad (1)$$

where $Z(\beta, B) = \sum_x e^{-\beta H(x) + B \sum_i x_i}$ is a normalising constant to make G a probability measure, β and B are constants representing the inverse temperature and external field respectively, and H is the *Hamiltonian*

$$H_{N,M}^{\text{Hop}}(x; \xi) = -\frac{1}{\sqrt{NM}} \sum_{k=1}^M (x \cdot \xi^k)^2 = -\frac{1}{\sqrt{N}} \sum_{i=1}^N \sum_{j=1}^N \left(\frac{1}{\sqrt{M}} \sum_{k=1}^M \xi_i^k \xi_j^k \right) x_i x_j. \quad (2)$$

The special configurations $\xi^k \in \mathbb{R}^N$, $1 \leq k \leq M$, are called *patterns* for the reason that the Gibbs measure tends to be concentrated near them, and thus these patterns are memorised in some sense (Hopfield, 1982). The patterns themselves are random, making the Gibbs measure a random measure; usually they are taken to be vectors of independent Bernoulli variables, but we will also consider more general patterns.

Our Hamiltonian is normalised by $1/\sqrt{NM}$ rather than the usual $1/N$. This is the correct normalisation as $\alpha = M/N \rightarrow \infty$; observe that if $1/\sqrt{M}$ were replaced by $1/\sqrt{N}$ in (2), then the expression in parentheses would diverge as $\alpha \rightarrow \infty$, resulting in a degenerate Gibbs measure. As a result of this normalisation, our inverse temperature β differs from the usual inverse temperature by a factor of $\sqrt{\alpha}$.

An important quantity associated with the Gibbs measure is the *free energy*

$$F(\beta, B) = \frac{1}{N} \log Z(\beta, B) = \frac{1}{N} \log \sum_x e^{-\beta H(x) + B \sum_i x_i}. \quad (3)$$

The significance of the free energy is that expectations with respect to the Gibbs measure contain a factor of $1/Z$, so many useful quantities appear as derivatives of F .

A related model is the Sherrington-Kirkpatrick (SK) model, with Hamiltonian

$$H_N^{SK}(x; J) = -\frac{1}{\sqrt{N}} \sum_{i=1}^N \sum_{j=1}^N J_{ij} x_i x_j. \quad (4)$$

The J_{ij} represent ferromagnetic *interactions* between sites i and j , and are usually taken to be independent standard Gaussian random variables. In this case, [Guerra and Toninelli \(2002\)](#) proved that the SK free energy converges to a limit (with exponentially high probability). The value of the limit was conjectured by [Parisi \(1980\)](#) and proved by [Guerra \(2003\)](#) and [Talagrand \(2006\)](#), although it is difficult to compute. [Carmona and Hu \(2003\)](#) showed that the limit holds for any interactions which are independent (not necessarily identically distributed) with mean 0, variance 1 and bounded third moment.

Our main result is that the Hopfield free energy, when appropriately normalised, converges to this same limit as the ratio $\alpha = M/N$ of patterns to sites grows large. We require the patterns to be symmetric, independent and identically distributed, with unit variance and bounded eleventh moment. Some generalisations which are not difficult to prove include replacing the symmetric requirement with normal concentration, removing the identically distributed requirement or including more general external fields, but we will not discuss them further.

Theorem 1. *If $\alpha \geq 5\beta^2$, then for some universal constant C ,*

$$\left| \mathbb{E}[F_{N,M}^{Hop}(\beta, B; \xi)] - \mathbb{E}[F_N^{SK}(\sqrt{2}\beta, B; J)] - \beta\sqrt{\alpha} \right| \leq \frac{C\beta^3}{\sqrt{\alpha}} + O\left(\frac{\beta^3}{\sqrt{M}}\right). \quad (5)$$

It is known ([Talagrand, 1996](#)) that the Hopfield free energy concentrates for Bernoulli patterns as $N \rightarrow \infty$; we give a proof for more general patterns in [Theorem 8](#). This gives the following behaviour of the limiting free energy with respect to α , which is confirmed in [Figure 1](#).

Corollary 2. *Let $P(\beta, B) = \lim_{N \rightarrow \infty} F_N^{SK}(\beta, B; J)$ be the limiting SK free energy. Then,*

$$\lim_{N \rightarrow \infty} F_{N,\alpha N}^{Hop}(\beta, B; \xi) = \beta\sqrt{\alpha} + P(\sqrt{2}\beta, B) + O(\beta^3/\sqrt{\alpha}). \quad (6)$$

The Hopfield model with large α is not well-studied. The region $\alpha \rightarrow 0$ is the best understood ([Shcherbina and Tirozzi, 1993](#); [Bovier and Gayraud, 1997](#)), and the problem becomes considerably more difficult as soon as $\limsup \alpha > 0$ ([Bovier, Gayraud and Picco, 1995](#); [Talagrand, 1998, 2000](#)). The author is not aware of any known results for unbounded α .

Heuristically, one can guess that the Hopfield model behaves like the SK model for large α , since the interaction between sites i and j is $\frac{1}{\sqrt{M}} \sum_k \xi_i^k \xi_j^k$, which converges to Gaussian when the limits $\alpha \rightarrow \infty$ and $N \rightarrow \infty$ are swapped. However, the Hopfield interactions are certainly not independent, nor is it clear that we can exchange these limits.

[Theorem 1](#) shows that the order of the limits does not matter, while [Lemma 5](#) shows that the dependence between interactions exactly corresponds to the $\beta\sqrt{\alpha}$ term in [Theorem 1](#). In fact, this term comes entirely from self-interactions when $i = j$; if we restrict our Hamiltonians to $i < j$, then $F^{Hop}(\beta, B) \sim F^{SK}(\beta, B)$ with no $\beta\sqrt{\alpha}$ term. Note that the $\sqrt{2}$ factor in the inverse temperature also disappears since it arises from the fact that the interactions J_{ij} and J_{ji} are independent in the SK model but equal in the Hopfield model.

Our paper is similar in structure to the universality result of [Carmona and Hu \(2003\)](#) for the SK model, but their bounds do not carry over directly to the Hopfield model. The problem is that the Hopfield Hamiltonian is quadratic in the patterns while the SK Hamiltonian is linear in the interactions. We solve this problem by bounding various quantities of interest by moments of the *overlap* $S = (x \cdot \xi^1)/\sqrt{N}$, which are evaluated in Theorem 4 using the symmetry trick of Lemma 3. This is similar to Theorem 2.2 in [Talagrand \(2000\)](#), but applies to a different parameter region and has a much more elementary proof.




Figure 1. Plot of free energy (vertical axis) against the parameter α (horizontal axis), for $\beta = 1$, $B = 0$ (bottom plot) and $\beta = 2$, $B = 5$ (top plot). The points are realisations of the Hopfield free energy with $N = 50$ and $\alpha = 1, 2, \dots, 50$. The curves are $\beta\sqrt{\alpha} + P(\sqrt{2}\beta, B)$, estimated by averaging 100 realisations of $F_{50}^{SK}(\sqrt{2}\beta, B)$.

2 Main Result

For $0 \leq t \leq 1$, define the interpolated Hamiltonian

$$H_{N,M}^t(x; \xi, J) = \sqrt{1-t}\sqrt{2}H_N^{SK}(x; J) + \sqrt{t}H_{N,M}^{Hop}(x; \xi) + N\sqrt{\alpha t} \quad (7)$$

$$= - \sum_{i,j} \left(\frac{\sqrt{1-t}\sqrt{2}}{\sqrt{N}} J_{ij} + \frac{\sqrt{t}}{\sqrt{NM}} \sum_k \xi_i^k \xi_j^k \right) x_i x_j + N\sqrt{\alpha t}. \quad (8)$$

Also define the corresponding partition function $Z = \sum_x e^{-\beta H(x) + B \sum_i x_i}$, Gibbs measure $G(x) = e^{-\beta H(x) + B \sum_i x_i} / Z$, and free energy $F = \frac{1}{N} \log Z$. Let $\langle f \rangle = \sum_x G(x) f(x)$ denote expectation with respect to the Gibbs measure. For convenience, we will often drop implied dependences; we refer to the interpolated quantities unless explicitly stated otherwise.

Assume the SK interactions J are independent standard Gaussian, and the Hopfield patterns ξ are symmetric, independent and identically distributed with unit variance; we will state more assumptions when they are required. We begin by giving a bound on moments of the overlap $S = (x \cdot \xi^1) / \sqrt{N} = \frac{1}{\sqrt{N}} \sum_i x_i \xi_i^1$, which will be crucial to the remainder of the paper.

Lemma 3. *Suppose $\sqrt{\alpha} \geq (2+3\epsilon)\beta$ for some $\epsilon > 0$, and let $A = \{\|\xi^1\|_2^2 > (1+\epsilon)N\}$. Then, for $c > 0$ and $C < \infty$ depending only on ϵ , and all $r > 0$,*

$$\mathbb{E}[G(S^2 > r) \mathbb{1}_{A^c}] \leq C e^{-cr}. \quad (9)$$

Proof. Let $E_u = \{u < S^2 \leq u+1\}$, considered as either a set of configurations or a set of patterns depending on context. Let $H^* = H + \sqrt{t}S^2/\sqrt{\alpha}$ be the Hamiltonian with the ξ^1 term removed, and let G^* be the corresponding Gibbs measure with expectation $\langle \cdot \rangle^*$. Then,

$$G(S^2 > r) = \sum_{u=r}^{\infty} G(E_u) = \sum_{u=r}^{\infty} \frac{\sum_{x \in E_u} e^{\beta \sqrt{t} S^2 / \sqrt{\alpha} - \beta H^*(x) + B \sum_i x_i}}{\sum_x e^{\beta \sqrt{t} S^2 / \sqrt{\alpha} - \beta H^*(x) + B \sum_i x_i}} \quad (10)$$

$$\leq \sum_{u=r}^{\infty} e^{\beta(u+1)/\sqrt{\alpha}} \frac{\sum_{x \in E_u} e^{-\beta H^*(x) + B \sum_i x_i}}{\sum_x e^{-\beta H^*(x) + B \sum_i x_i}} = \sum_{u=r}^{\infty} e^{\beta(u+1)/\sqrt{\alpha}} G^*(E_u). \quad (11)$$

Since G^* does not depend on ξ^1 , by Tonelli's theorem, we can exchange $\mathbb{E}_{\xi^1}$ and $\langle \cdot \rangle^*$ to obtain

$$\mathbb{E}[G(S^2 > r) \mathbb{1}_{A^c}] \leq \sum_{u=r}^{\infty} e^{\beta(u+1)/\sqrt{\alpha}} \mathbb{E}\left[\left\langle \mathbb{P}_{\xi^1}(E_u \cap A^c) \right\rangle^*\right]. \quad (12)$$

Since replacing ξ_i^1 with $x_i \xi_i^1$ changes neither the distribution of S nor the value of $\|\xi^1\|_2^2$, $\mathbb{P}_{\xi^1}(E_u \cap A^c)$ does not depend on x . Then, by [Hoeffding's \(1963\)](#) inequality,

$$\mathbb{P}_{\xi^1}(E_u \cap A^c) = \frac{1}{2^N} \sum_x \mathbb{P}_{\xi^1}(E_u \cap A^c) = \mathbb{E}\left[\frac{|E_u|}{2^N} \mathbb{1}_{A^c}\right] \leq \mathbb{E}\left[2e^{-uN/2\|\xi^1\|_2^2} \mathbb{1}_{A^c}\right]. \quad (13)$$

Since $A^c = \{\|\xi^1\|_2^2 \leq (1+\epsilon)N\}$, this is bounded by $2e^{-u/(2+2\epsilon)}$, hence

$$\mathbb{E}[G(S^2 > r) \mathbb{1}_{A^c}] \leq 2 \sum_{u=r}^{\infty} \exp\left(\frac{\beta(u+1)}{\sqrt{\alpha}} - \frac{u}{2+2\epsilon}\right) \quad (14)$$

Our assumption $\sqrt{\alpha} \geq (2+3\epsilon)\beta$ implies the required bound. $\square$

Theorem 4. Suppose $\sqrt{\alpha} \geq (2+3\epsilon)\beta$, and the patterns ξ have bounded m th moment for some $m > 2$. For any d such that $0 \leq d < (m-2)(1-d/m)$, there is a constant C_d depending only on ϵ and $\mathbb{E}[|\xi_1^1|^m]$ such that $\mathbb{E}[\langle |S|^d \rangle] \leq C_d$.

Proof. Let $A = \{ \|\xi^1\|_2^2 > (1+\epsilon)N \}$. Then,

$$\mathbb{E}[\langle |S|^d \rangle] = \mathbb{E}[\langle |S|^d \rangle \mathbb{1}_A] + \mathbb{E}[\langle |S|^d \rangle \mathbb{1}_{A^c}] \quad (15)$$

$$\leq \mathbb{E}[\langle |S|^m \rangle]^{d/m} \mathbb{P}[A]^{1-d/m} + \int_0^\infty \mathbb{E}[G(S^2 > r^{2/d}) \mathbb{1}_{A^c}] dr. \quad (16)$$

By Lemma 3, this integral is bounded. Using the trivial bound $\mathbb{E}[\langle |S|^m \rangle] \leq N^{m/2} \mathbb{E}[|\xi_1^1|^m]$, since $(m-2)(1-d/m) > d$, it suffices to prove $\mathbb{P}[A] = O(N^{-(m-2)/2}(\log N)^m)$.

Let $f(r) = r^m \mathbb{P}[|\xi_1^1| > r]$. We claim that there exists c such that for any $r > 0$, the interval $[r, cr]$ contains a point u with $f(u) < 1$. We prove this by contradiction; if the claim is false, then for any c , there is r_c such that $f \geq 1$ on $[r_c, cr_c]$, and we arrive at the contradiction

$$\mathbb{E}[|\xi_1^1|^m] = \int_0^\infty r^{m-1} \mathbb{P}[|\xi_1^1| > r] dr = \int_0^\infty \frac{f(r)}{r} dr \geq \int_{r_c}^{cr_c} \frac{1}{r} dr = \log c. \quad (17)$$

Thus, we can pick r_N such that $\sqrt{N}/\log N \leq r_N \leq c\sqrt{N}/\log N$ and $\mathbb{P}[|\xi_1^1| > r_N] < r_N^{-m}$. Let $B = \{\forall i |\xi_i^1| \leq r_N\}$. Then,

$$\mathbb{P}[B^c] \leq N \mathbb{P}[|\xi_1^1| > r_N] \leq N r_N^{-m} \leq N^{1-m/2} (\log N)^m. \quad (18)$$

Finally, by Hoeffding's (1963) inequality, $\mathbb{P}[B \cap A] \leq 2e^{-\epsilon N/2r_N^2} \leq 2e^{-\epsilon(\log N)^2/2c^2}$. $\square$

The next two lemmas are formulae for evaluating quantities that will be important later, while Lemma 7 is a universality result establishing that the distribution of patterns does not matter, thus allowing us to apply Stein's lemma as if they were Gaussian.

Lemma 5. The interpolated mean free energy has derivative

$$\frac{d}{dt} \mathbb{E}[F] = \frac{\beta(N-1)}{\sqrt{2(1-t)N}} \mathbb{E}[J_{12} \langle x_1 x_2 \rangle] - \frac{\beta(N-1)M}{2\sqrt{tNM}} \mathbb{E}[\xi_1^1 \xi_2^1 \langle x_1 x_2 \rangle]. \quad (19)$$

Proof. We have

$$\frac{d}{dt} \mathbb{E}[F] = \mathbb{E} \left[\frac{1}{NZ} \sum_x \left(-\beta \frac{\partial H}{\partial t} \right) e^{-\beta H(x) + B \sum_i x_i} \right] = -\frac{\beta}{N} \mathbb{E} \left[\left\langle \frac{\partial H}{\partial t} \right\rangle \right]. \quad (20)$$

Next, we differentiate (8) and split the sum over i and j into two sums where $i \neq j$ and $i = j$ respectively. The $i \neq j$ terms give (19) upon taking $i = 1$, $j = 2$ and $k = 1$ without loss of generality. For the $i = j$ terms, the SK term is a multiple of $\mathbb{E}[J_{11} \langle x_1^2 \rangle] = 0$, while the Hopfield term exactly cancels the derivative of $N\sqrt{\alpha t}$. $\square$

Lemma 6. For any function $g = g(x; \xi, J)$, and $S = (x \cdot \xi^1)/\sqrt{N}$,

$$\frac{\partial}{\partial \xi_i^1} \langle g \rangle = \left\langle \frac{\partial g}{\partial \xi_i^1} \right\rangle + \frac{2\beta\sqrt{t}}{\sqrt{M}} \left(\langle g x_i S \rangle - \langle g \rangle \langle x_i S \rangle \right). \quad (21)$$

Proof. Explicit derivative calculation. $\square$

Lemma 7. Suppose $\alpha \geq (4 + \epsilon)\beta^2$ for some $\epsilon > 0$, and the patterns ξ have bounded m th moment for $m = 11$. Then, for some constant C depending only on ϵ and $\mathbb{E}[|\xi_1^1|^{11}]$,

$$\left| \mathbb{E}[\xi_1^1 \xi_2^1 \langle x_1 x_2 \rangle] - \mathbb{E}\left[\frac{\partial}{\partial \xi_1^1} \frac{\partial}{\partial \xi_2^1} \langle x_1 x_2 \rangle\right] \right| \leq \frac{C\beta^2}{M\sqrt{N}}. \quad (22)$$

Proof. Define the probabilistic function $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ by $f = \langle x_1 x_2 \rangle$ considered as a function of $y = \xi_1^1$ and $z = \xi_2^1$, with the other ξ_i^k regarded as random parameters. Since f is infinitely differentiable, we can define its second degree Taylor expansion T around 0, that is,

$$T(y, z) = f(0, 0) + yf_y(0, 0) + zf_z(0, 0) + \frac{1}{2}y^2 f_{yy}(0, 0) + \frac{1}{2}z^2 f_{zz}(0, 0) + yzf_{yz}(0, 0). \quad (23)$$

Since the evaluated derivatives depend only on the ξ_i^k other than y or z , which are independent with mean 0 and variance 1, an easy calculation yields $\mathbb{E}[yzT(y, z)] = \mathbb{E}[f_{yz}(0, 0)]$, hence

$$\mathbb{E}[yzf(y, z) - f_{yz}(y, z)] = \mathbb{E}[yz(f - T)(y, z) - (f_{yz}(y, z) - f_{yz}(0, 0))]. \quad (24)$$

By Taylor's theorem,

$$\begin{aligned} (f - T)(y, z) &= \int_0^1 \left(\frac{1}{2}y^3 f_{yyy}(sy, sz) + \frac{3}{2}y^2 z f_{yyz}(sy, sz) \right. \\ &\quad \left. + \frac{3}{2}yz^2 f_{yzz}(sy, sz) + \frac{1}{2}z^3 f_{zzz}(sy, sz) \right) (1-s)^2 ds; \end{aligned} \quad (25)$$

$$f_{yz}(y, z) - f_{yz}(0, 0) = \int_0^1 \left(y f_{yyz}(sy, sz) + z f_{yzz}(sy, sz) \right) ds. \quad (26)$$

By symmetry of $f(y, z) = f(z, y)$,

$$\begin{aligned} \mathbb{E}[yzf(y, z) - f_{yz}(y, z)] &= \int_0^1 (1-s^2) \mathbb{E}[y^4 z f_{yyy}(sy, sz)] \\ &\quad + \mathbb{E}[(3(1-s)^2 y^3 z^2 + 2y) f_{yyz}(sy, sz)] ds. \end{aligned} \quad (27)$$

Let $X = x_1 x_2$ and $U = x_1 S$. By Lemma 6,

$$f_y = \frac{\partial}{\partial \xi_1^1} \langle X \rangle = \frac{2\beta\sqrt{t}}{\sqrt{M}} (\langle XU \rangle - \langle X \rangle \langle U \rangle); \quad (28)$$

$$f_{yy} = \left(\frac{2\beta\sqrt{t}}{\sqrt{M}} \right)^2 (\langle XU^2 \rangle - 2\langle XU \rangle \langle U \rangle - \langle X \rangle \langle U^2 \rangle + 2\langle X \rangle \langle U \rangle^2); \quad (29)$$

$$\begin{aligned} f_{yyy} &= \left(\frac{2\beta\sqrt{t}}{\sqrt{M}} \right)^3 \left(\langle XU^3 \rangle - 3\langle XU^2 \rangle \langle U \rangle - 3\langle XU \rangle \langle U^2 \rangle + 6\langle XU \rangle \langle U \rangle^2 \right. \\ &\quad \left. - \langle X \rangle \langle U^3 \rangle + 6\langle X \rangle \langle U^2 \rangle \langle U \rangle - 6\langle X \rangle \langle U \rangle^3 \right). \end{aligned} \quad (30)$$

By Hölder's inequality, $|f_{yyy}| \leq 8\beta^3 M^{-3/2} 26 \langle |S|^3 \rangle$. Let $\langle \cdot \rangle^s$ be expectation with respect to the Gibbs measure with ξ_1^1 and ξ_2^1 replaced by $s\xi_1^1$ and $s\xi_2^1$ respectively. Then,

$$\left| \mathbb{E}[y^4 z f_{yyy}(sy, sz)] \right| \leq \frac{8\beta^3}{M^{3/2}} 26 \mathbb{E} \left[|y|^4 |z| \left\langle |S - (1-s)(x_1 y + x_2 z)|^3 \right\rangle^s \right]. \quad (31)$$

Thus, we need to bound $\mathbb{E}[\langle |y|^{4+a}|z|^{1+b}|S|^c \rangle^s]$, where $a+b+c=3$. By Hölder's inequality, this is bounded by $\mathbb{E}[|y|^{(4+a)d/(d-c)}|z|^{(1+b)d/(d-c)}]^{1-c/d} \mathbb{E}[\langle |S|^d \rangle]^{c/d}$, where $(4+a)d/(d-c) \leq m$ and $c < d < (m-2)(1-d/m)$. By Theorem 4, which applies to $\langle \cdot \rangle^s$ since scaling finitely many patterns does not affect the proof, this yields $\mathbb{E}[y^4 z f_{yyy}(sy, sz)] = O(\beta^3/M^{3/2})$; some algebra shows we need $m > 6 + \sqrt{22} \approx 10.7$.

Bounding the second expectation in (27) is similar. Using Lemma 6 to differentiate (29) with respect to z and applying Hölder's inequality gives

$$|f_{yyz}| \leq \frac{4\beta^2}{M\sqrt{N}} 8\langle |S| \rangle + \frac{8\beta^3}{M^{3/2}} 26\langle |S|^3 \rangle. \quad (32)$$

Scaling ξ_1^1 and ξ_2^1 by s , multiplying by $3(1-s)^2 y^3 z^2 + 2y$ and taking expectations, the same argument gives the bound $O(\beta^2/M\sqrt{N})$, noting that $\beta/\sqrt{\alpha} < \frac{1}{2}$. $\square$

Theorem 1. *Suppose the patterns ξ have bounded eleventh moment. If $\alpha \geq (4+\epsilon)\beta^2$ for some $\epsilon > 0$, then for some constant C depending only on ϵ ,*

$$\left| \mathbb{E}[F_{N,M}^{Hop}(\beta, B; \xi)] - \mathbb{E}[F_N^{SK}(\sqrt{2}\beta, B; J)] - \beta\sqrt{\alpha} \right| \leq \frac{C\beta^3}{\sqrt{\alpha}} + O\left(\frac{\beta^3}{\sqrt{M}}\right). \quad (33)$$

Proof. Observe that

$$F_{N,M}^{Hop}(\beta, B; \xi) - F_N^{SK}(\sqrt{2}\beta, B; J) - \beta\sqrt{\alpha} = \int_0^1 \frac{d}{dt} \mathbb{E}[F_{N,M}^t(\beta, B; \xi, J)] dt. \quad (34)$$

By Lemma 5, it suffices to prove

$$\left| \frac{\beta(N-1)}{\sqrt{2(1-t)N}} \mathbb{E}[J_{12}\langle x_1 x_2 \rangle] - \frac{\beta(N-1)M}{2\sqrt{tNM}} \mathbb{E}[\xi_1^1 \xi_2^1 \langle x_1 x_2 \rangle] \right| \leq \frac{C\beta^3}{\sqrt{\alpha}} + O\left(\frac{\beta^3}{\sqrt{M}}\right). \quad (35)$$

Let $X = x_1 x_2$ and $\text{Var}(X) = \langle X^2 \rangle - \langle X \rangle^2$. By Stein's lemma,

$$\mathbb{E}[J_{12}\langle x_1 x_2 \rangle] = \mathbb{E}\left[\frac{\partial}{\partial J_{12}} \frac{\sum_x x_1 x_2 e^{-\beta H(x) + B \sum_i x_i}}{\sum_x e^{-\beta H(x) + B \sum_i x_i}} \right] = \frac{\beta \sqrt{2(1-t)}}{\sqrt{N}} \mathbb{E}[\text{Var}(X)]. \quad (36)$$

Let $U = x_1 S$ and $V = x_2 S$. By Lemmas 6 and 7 applied to (28),

$$\begin{aligned} \mathbb{E}[\xi_1^1 \xi_2^1 \langle x_1 x_2 \rangle] &= \frac{4\beta^2 t}{M} \mathbb{E}[\langle XUV \rangle - \langle XU \rangle \langle V \rangle - \langle XV \rangle \langle U \rangle - \langle X \rangle \langle UV \rangle + 2\langle X \rangle \langle U \rangle \langle V \rangle] \\ &\quad + \frac{2\beta\sqrt{t}}{\sqrt{NM}} \mathbb{E}[\text{Var}(X)] + O\left(\frac{\beta^2}{M\sqrt{N}}\right). \end{aligned} \quad (37)$$

By Hölder's inequality and Theorem 4, the first line is $O(\beta^2/M)$, which gives the $C\beta^3/\sqrt{\alpha}$ term. In the second line, the first term exactly cancels (36), while the second term gives the $O(\beta^3/\sqrt{M})$ error. It is clear that C depends only on ϵ and $\mathbb{E}[|\xi_1^1|^{11}]$; the moment dependence only arises in the $N^{-\log N}$ term in Theorem 4 and does not affect the dominant term. $\square$

Finally, we prove the concentration that allows us to conclude Corollary 2 from Theorem 1.

Theorem 8. *Suppose $\alpha \geq (4 + \epsilon)\beta^2$ for some $\epsilon > 0$, and the patterns ξ have bounded m th moment. If $4 \leq 2d < (m - 2)(1 - 2d/m)$, then for some C depending only on ϵ and $\mathbb{E}[|\xi_1^1|^m]$,*

$$\mathbb{E} \left[\left| F_{N, \alpha N}^{\text{Hop}}(\beta, B; \xi) - \mathbb{E}[F_{N, \alpha N}^{\text{Hop}}(\beta, B; \xi)] \right|^d \right] \leq \frac{C\beta^d}{N^{d/2}}. \quad (38)$$

Proof. We follow the proof of Carmona and Hu (2003) for the SK model, using our bound from Theorem 4. Since we are only dealing with the Hopfield free energy here, symbols in this proof will refer to the Hopfield quantities rather than the interpolated ones.

Let $\mathcal{F}_\ell$ be the σ -algebra generated by $\xi^1, \dots, \xi^\ell$, and let $D_\ell = N(\mathbb{E}[\log Z | \mathcal{F}_\ell] - \mathbb{E}[\log Z | \mathcal{F}_{\ell-1}])$. The D_ℓ are martingale differences with $\sum_\ell D_\ell = N(F - \mathbb{E}[F])$, so by Burkholder's (1973) martingale inequality and convexity of $x \mapsto x^{d/2}$, for some universal constant C_d ,

$$\mathbb{E}[|F - \mathbb{E}[F]|^d] = \frac{1}{N^d} \mathbb{E} \left[\left| \sum_\ell D_\ell \right|^d \right] \leq \frac{C_d}{N^d} \mathbb{E} \left[\left(\sum_\ell D_\ell^2 \right)^{d/2} \right] \leq \frac{C_d M^{d/2}}{N^d} \frac{1}{M} \sum_\ell \mathbb{E}[|D_\ell|^d]. \quad (39)$$

Let $\beta' = \beta/\sqrt{NM}$. Observe that for any fixed $1 \leq \ell \leq M$, we can write

$$\log Z = \log \sum_x e^{\beta' \sum_{k \neq \ell} (x \cdot \xi^k)^2 + B \sum_i x_i} - \log \frac{\sum_x e^{\beta' \sum_{k \neq \ell} (x \cdot \xi^k)^2 + B \sum_i x_i}}{\sum_x e^{\beta' \sum_k (x \cdot \xi^k)^2 + B \sum_i x_i}} \quad (40)$$

$$= \log \sum_x e^{\beta' \sum_{k \neq \ell} (x \cdot \xi^k)^2 + B \sum_i x_i} - \log \left\langle e^{-\beta' (x \cdot \xi^\ell)^2} \right\rangle. \quad (41)$$

Since the first term above is independent of ξ^ℓ , it follows that

$$D_\ell = -\mathbb{E} \left[\log \left\langle e^{-\beta' (x \cdot \xi^\ell)^2} \right\rangle \middle| \mathcal{F}_\ell \right] + \mathbb{E} \left[\log \left\langle e^{-\beta' (x \cdot \xi^\ell)^2} \right\rangle \middle| \mathcal{F}_{\ell-1} \right]. \quad (42)$$

By Hölder's inequality,

$$\mathbb{E}[|D_\ell|^d] \leq 2^d \mathbb{E} \left[\left| \log \left\langle e^{-\beta' (x \cdot \xi^\ell)^2} \right\rangle \right|^d \right] = 2^d \mathbb{E} \left[\left| \log \left\langle e^{-\beta S^2 / \sqrt{\alpha}} \right\rangle \right|^d \right]. \quad (43)$$

The function $\phi(z) = |\log z|^d$ has second derivative $\phi''(z) = (d(d-1) |\log z|^{d-2} - d |\log z|^{d-1})/z^2$, so ϕ is convex for $0 < z \leq e^{d-1}$. Since $0 < e^{-\beta' S^2} \leq 1 < e^{d-1}$, by Jensen's inequality,

$$\mathbb{E}[|D_\ell|^d] \leq 2^d \mathbb{E} \left[\left\langle \left| \log (e^{-\beta S^2 / \sqrt{\alpha}}) \right|^d \right\rangle \right] = 2^d \mathbb{E} \left[\left\langle \left(\frac{\beta S^2}{\sqrt{\alpha}} \right)^d \right\rangle \right]. \quad (44)$$

This is $O(\beta^d / \alpha^{d/2})$ by Theorem 4. $\square$

Acknowledgements. The author wishes to thank Amir Dembo, Persi Diaconis, Jack Kamm and Andrea Montanari for their helpful suggestions which contributed greatly to this paper.

References

- A. Bovier and V. Gayrard. An almost sure central limit theorem for the Hopfield model. *Markov Processes and Related Fields* **3**(2):151–173, 1997. [MR1468172](#).
- A. Bovier and V. Gayrard. Hopfield models as generalized random mean field models. In A. Bovier and P. Picco, editors, *Mathematical Aspects of Spin Glasses and Neural Networks*, pages 3–90. Birkhäuser, Boston, 1997.
- A. Bovier, V. Gayrard and P. Picco. Gibbs states of the Hopfield model with extensively many patterns. *Journal of Statistical Physics* **79**(1–2):395–414, 1995. [MR1325589](#).
- D.L. Burkholder. Distribution function inequalities for martingales. *Annals of Probability* **1**(1):19–42, 1973. [MR0365692](#).
- P. Carmona and Y. Hu. Universality in Sherrington-Kirkpatrick’s spin glass model. *Annales de l’Institut Henri Poincaré* **42**(2):215–222, 2006. [MR2199799](#).
- F. Guerra. Broken replica symmetry bounds in the mean field spin glass model. *Communications in Mathematical Physics* **233**(1):1–12, 2003. [MR1957729](#).
- F. Guerra and L.T. Toninelli. The thermodynamic limit in mean field spin glass models. *Communications in Mathematical Physics* **230**(1):71–79, 2002. [MR1930572](#).
- W. Hoeffding. Probability inequalities for sums of bounded random variables. *Journal of the American Statistical Association* **58**(301):13–30, 1963. [MR0144363](#).
- J.J. Hopfield. Neural networks and physical systems with emergent collective computational abilities. *Proceedings of the National Academy of Sciences of the USA* **79**(8):2554–2558, 1982. [MR0652033](#).
- M. Ledoux. *The Concentration of Measure Phenomenon*. American Mathematical Society, Providence, 2001.
- G. Parisi. A sequence of approximated solutions to the S-K model for spin glasses. *Journal of Physics A: Mathematical and General* **13**(4):L115–L121, 1980.
- M. Shcherbina and B. Tirozzi. The free energy of a class of Hopfield models. *Journal of Statistical Physics* **72**(1–2):113–125, 1993. [MR1233028](#).
- M. Talagrand. A new look at independence. *Annals of Probability* **24**(1):1–34, 1996. [MR1387624](#).
- M. Talagrand. Rigorous results for the Hopfield model with many patterns. *Probability Theory and Related Fields* **110**(2):177–276, 1998. [MR1609015](#).
- M. Talagrand. Exponential inequalities and convergence of moments in the replica-symmetric regime of the Hopfield model. *Annals of Probability* **28**(4):1393–1469, 2000. [MR1813831](#).
- M. Talagrand. The Parisi formula. *Annals of Mathematics* **163**(1) 221–263, 2006. [MR2195134](#).